

Differences in mental health problems, coping self-efficacy and social support between adults victimised before and adults victimised after the COVID-19 outbreak: population-based prospective study

Peter G. van der Velden, Carlo Contino, Marcel Das, Joost Leenen and Lutz Wittmann

Background

Victims of violence, accidents and threats are at risk for mental health problems. Lower coping self-efficacy and social support levels increase this risk. Although highly relevant, it is unknown if the coronavirus disease 2019 (COVID-19) pandemic amplifies these risks.

Aims

To examine if the prevalence, incidence and/or mean scores for post-traumatic stress disorder (PTSD), anxiety and depression symptoms, general mental health problems, coping self-efficacy, lack of emotional support and social acknowledgement are higher among adults victimised in the year after the COVID-19 outbreak compared with adults victimised in a similar period before the outbreak. Also, to compare symptoms, problems and support within non-victims during the same period.

Method

Data was extracted from four surveys of the VICTIMS study (March 2018, 2019, 2020, 2021), based on a random sample of the Dutch population. Multivariate logistic regression analyses and mixed-effects models were used to examine differences between the two victim groups (2019: $n = 421$, 2021: $n = 319$) and non-victims ($n = 3245$).

Results

Adults victimised after the outbreak more often had PTSD, anxiety and depression symptoms, general mental health problems

and lower coping self-efficacy than those victimised before. They did not differ in lack of support and acknowledgement. Both victim groups differed from non-victims, where mental health problems and lack of support levels were much lower and almost stable.

Conclusions

The COVID-19 pandemic had a negative impact on the mental health and coping self-efficacy levels of victims, whereas mental health problems among non-victims remained virtually stable. Mental healthcare workers, general practitioners and victim services should take this impact into account.

Keywords

COVID-19; PTSD; social support; coping self-efficacy; victims.

Copyright and usage

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Background

Each year, a considerable minority of the adult general population is victimised by exposure to violence, accidents and serious threats (VATs).¹ The effects of these events on the mental health of the victims are well documented.² With respect to risk factors for post-trauma mental health problems, research has shown that lower levels of event-related coping self-efficacy, for example, the perceived ability to cope with post-trauma recovery demands^{3,4} and lower levels of social support are associated with an increased risk of post-trauma mental health problems.^{5,6}

On 11 March 2020, the coronavirus disease 2019 (COVID-19) outbreak was officially declared a pandemic by the World Health Organization. A meta-analysis of longitudinal cohort studies on the effects of this pandemic on the mental health of the general population with pre-outbreak data on mental health, has shown that the overall increase in mental health symptoms was most pronounced during the early stages of the pandemic (March–April 2020), before decreasing and being generally comparable with pre-pandemic levels by mid-2020.⁷ However, even in the absence of enduring effects of the current pandemic on the mental health of the general population, specific subgroups may be less resilient to pandemic-

related stressors. As it is known that psychosocial problems such as financial hardship or lack of social support can increase the impact of potentially traumatic events, ongoing COVID-19 pandemic-related stressors (for example lockdowns, infection and health risks, unemployment risk) may substantially disturb or hinder the normal recovery process of victims.^{8,9} Yet, to the best of our knowledge, to date no study has compared adults victimised during the COVID-19 pandemic with adults victimised in a similar time span before this pandemic. Insight into these effects is relevant for mental healthcare professionals and policymakers.

Aims

To fill this gap, the present population-based study compared the mental health (anxiety and depression symptoms, post-traumatic stress disorder (PTSD) symptoms, general mental health problems), event-related coping self-efficacy and lack of social support (lack of emotional support and social acknowledgement) of adults victimised by VATs before the outbreak (group 1^{victims}: victimised between March 2018 (T_1) and March 2019 (T_2)) with adults victimised by VATs in the year after the outbreak of COVID-19

(group 2^{victims}: victimised between March 2020 (T_3) and March 2021 (T_4)).

To examine the extent to which this pandemic specifically affects victims, we also examined the patterns of anxiety and depression symptoms, general mental health problems, and lack of emotional support among respondents in March 2019 and March 2021 not victimised by VATs between March 2018 and March 2021 (i.e. non-victims).

Method

Procedures and participants

For the present study, data was extracted from four waves of the prospective VICTIMS study^{1,6} conducted in March 2018 (T_1 ; $n^{\text{invited}} = 7292$, response 82.1%), March 2019 (T_2 ; $n^{\text{invited}} = 6298$, response 83.2%), March 2020 (T_3 ; $n^{\text{invited}} = 6568$, response 83.6%) and March 2021 (T_4 ; $n^{\text{invited}} = 6452$, response 86.7%).

The VICTIMS study was conducted using the Longitudinal Internet Studies for the Social Sciences (LISS) panel, based on a traditional probability sample drawn from the Dutch population register of 16 years and older by Statistics Netherlands. The set-up of LISS was funded by the Dutch Research Council (NWO) and managed by Centerdata.¹⁰ Panel members complete online questionnaires every month, which take about 30 min in total to complete. They receive an incentive of 15 euros per h and members who do not have a computer and/or internet access are provided with the necessary equipment at home (for further information about the LISS panel and open access data see: <https://www.dataarchive.lissdata.nl>, in English). The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures were approved by an Internal Review Board of Centerdata. Explicit digital informed consent was obtained from all respondents.

Victims

We first compared the two victim groups: respondents victimised between March 2018 and March 2019 (group 1^{victims}) and respondents victimised between March 2020 and March 2021 (group 2^{victims}). For group 1^{victims}, we linked data on mental health and lack of support assessed at T_1 (before victimisation between T_1 and T_2) and for group 2^{victims} we linked the same data assessed at T_3 (before victimisation between T_3 and T_4).

Preliminary analyses showed that 105 respondents were victimised between T_1 and T_2 , as well as between T_3 and T_4 , prohibiting an independent comparison between both victim groups. To solve this problem, we randomly divided this subgroup into two subgroups. One-half ($n = 52$) was added to the group of respondents who participated at T_1 and T_2 (group 1^{total}, $n = 4717$) and the other half ($n = 53$) to respondents who participated at T_3 and T_4 (group 2^{total}, $n = 4875$). In this way, the proportion of group 1^{victims} and group 2^{victims} is similar (56% versus 44%) to the original proportion of people with VAT cases in between T_1 and T_2 , and in between T_3 and T_4 .

Next, we weighted the data of group 1^{total} and group 2^{total} using 16 exclusive demographic profiles among the total adult Dutch population (2018 and 2020 respectively), based on the data of Statistics Netherlands (see: <https://opendata.cbs.nl/#/CBS/en/>). These profiles were constructed using the variables gender (male, female), age (18–34, 35–49, 50–64, 65 years and older) and marital status (married and unmarried). After this procedure and weighting, we finally selected the victims of group 1 ($n = 421$) and the (non-overlapping) victims of group 2 ($n = 319$).

Non-victims

To examine differences in mental health and social support among non-victims before and after the outbreak, we assembled a second data-set. For this data-set, we first selected respondents who participated at T_2 , T_3 and T_4 (regardless of being victimised, $n = 4020$). After this selection, we weighted the data using the 16 demographic profiles of the Dutch adult population (2019). We finally selected respondents who were not victimised by VATs in the period T_1 to T_4 ($n^{\text{non-victims}} = 3245$).

All results are based on weighted data (except Cronbach's alpha).

Measures

At each survey, gender, age, marital and employment status and education level are assessed. Questions about post-traumatic stress symptoms, coping self-efficacy and social acknowledgement were only administered to victims.

Experiences with VATs

At each survey, respondents were asked to indicate for each of the 21 listed potentially traumatic events, if they were exposed to this specific event in the past 12 months (1, no; 2, yes) including the following VAT events:

- physical violence (sexual violence/sexual abuse (not online); online sexual violence/sexual abuse; robbery; physical violence, but not by own partner; physical violence by own partner);
- accidents (traffic accidents, disasters, fire, medical errors); and
- threats (serious threat; without the use of physical violence (not online); online serious threat, without use of physical violence).

The list also included events like (sudden) death of a (significant) other, (online) theft and serious illness categorised here as other stressful life events (SLEs). Participants were finally given the opportunity to report events in the previous 12 months that were not listed, which were recoded afterwards into existing or new categories.¹

Lastly, they were asked when the (most drastic or stressful) event occurred (coded as 1 for 1 week ago to 8 for 7–12 months ago), and asked to rate the level of stress during the event (coded 1 for none to 5 for very much). To reduce the number of cells with low cell counts, both time since the event and stress during the event were dichotomised (time: 1, up to 2 months ago (1, 2, 3, 4 and 5); 2, 3–12 months ago (6, 7 and 8); stress: 1, not very stressful (1, 2 and 3); 2, very stressful (4 and 5)).

Anxiety and depression symptoms

In all surveys, anxiety and depression symptoms were assessed using the five-item Mental Health Inventory: this scale covers symptoms during the past month on six-point Likert scales, such as 'This past month I felt very anxious' and 'I felt depressed and gloomy' (0, never to 5, continuously).^{11,12} After recoding the negative formulated items, the total scores were computed and multiplied by four (range 0 to 100), where lower scores indicate higher anxiety and depression symptom levels (all Cronbach's alpha of group 1^{victims}, group 2^{victims} and non-victims ≥ 0.85). A cut-off score of ≤ 59 was used to identify respondents with moderate–severe anxiety and depression symptom levels.¹³

Post-traumatic stress symptoms

To examine PTSD symptomatology at T_2 and T_4 , victims were asked to fill in the eight-item version of PTSD Checklist for DSM-5 (PCL-5)^{14,15} with the VAT event (reported in the same survey)

in mind. When respondents reported two or more events, they were asked to focus on the most impactful or stressful VAT event when filling in the PCL-5. This event was presented on the screen. Examples of items are 'Feeling very upset when something reminded you of the stressful experience' and 'Loss of interest in activities that you used to enjoy'. The items of the PCL-5 have a five-point Likert scale (0, not at all to 4, extremely) and examine symptoms across the four symptom clusters of PTSD according to DSM-5 and focus on symptoms in the past month (range 0 to 32). Higher scores reflect higher PTSD-symptom levels (Cronbach's alpha of group 1^{victims} and group 2^{victims} ≥ 0.92). To identify victims with probable PTSD, a cut-off of ≥ 13 was applied.¹⁶

General mental health problems and physical problems

The brief screening Problems and Help Inventarisation List¹⁷ was administered at all surveys to examine the prevalence of various problems such as general mental health problems, financial problems and physical problems. All respondents were asked to indicate for each of the problem types listed whether they experience these problems or not (1, yes; 2, no). For the present study, data was extracted from the items 'psychological problems' (as a proxy for general mental health problems), and 'physical problems'.

Coping self-efficacy

To examine VAT-related coping self-efficacy among victims at T_2 and T_4 , the seven-item Coping Self-Efficacy (CSE-7)¹⁸ measure was administered with items such as 'dealing with frightening images or dreams about the event' and 'being optimistic since the event'. The CSE-7 was administered in the same way as the PCL-5. For each item, respondents rated their perceived self-efficacy level on a seven-point Likert scale (1, I am completely incapable of to 7, I am perfectly capable of; Cronbach's alpha of group 1^{victims} and group 2^{victims} ≥ 0.93). Higher scores reflect higher coping self-efficacy levels (range 7–49). A cut-off of ≤ 28 , corresponding to (about) 10% lowest scores, was used to identify victims with relatively low versus high coping self-efficacy levels.

Social acknowledgement

To assess event-related social acknowledgement among victims at T_2 and T_4 , the five-item general disapproval scale of the Social Acknowledgment Questionnaire (SAQ)¹⁹ was administered, in the same way as the PCL-5 and CSE-7. The items have five-point answer categories (1, totally disagree to 5, totally agree). Examples of items are 'There is not enough sympathy for what happened to me' and 'Most people cannot understand what I went through' (both Cronbach's alpha of group 1^{victims} and group 2^{victims} ≥ 0.85). Higher scores reflect lower social acknowledgement levels (range 5–25). A cut-off of ≥ 17 , corresponding to (about) the 10% highest scores, was used to identify respondents with a high lack of social acknowledgement.

Lack of emotional support

In all surveys, lack of emotional support in response to problems was examined among all respondents with the eight-item subscale lack of emotional support of the Social Support List-Discrepancy (SSL-D).^{20,21} Examples of items are 'comfort you' and 'help you to clarify your problems'. The SSL-D items apply four-point Likert scales (1, I miss this, I would like it to happen more often to 4, It happens too often). Low scores reflect a higher lack of emotional support levels (range 8–32) (Cronbach's alpha of group 1^{victims}, group 2^{victims} and non-victims ≥ 0.88). A cut-off of ≤ 16 , corresponding to (about) the 10% lowest scores, was used to identify respondents with a high lack of emotional support.

Statistical analyses

Differences in characteristics between both victim groups were assessed using a chi-square test. Of the non-VAT-related characteristics of the non-victims at T_2 , T_3 and T_4 , we only examined differences in exposure to other events (SLEs) in the past 12 months and physical problems because age, education level, marital status and employment status follow a natural pattern (slight change) over a period of 3 years.

Post-event differences between both victim groups

To examine differences between both victim groups for the prevalence of post-event (for group 1^{victims} at T_2 , and for group 2^{victims} at T_4) anxiety and depression symptoms, post-traumatic stress symptoms, general mental health problems, coping self-efficacy, lack of emotional support and social acknowledgement, multivariate logistic regression analyses were conducted. To test differences between both victim groups in a rigorous way and to rule out alternative explanations for differences between the two groups as much as possible, we controlled for characteristics that differed between both groups not only at a $P < 0.05$ level but also at a $P \leq 0.10$ level (see Table 1), for example gender, number of different VAT events in the past 12 months and general mental health problems in the year before.

To examine differences in mean scores for post-event anxiety and depression symptoms, post-traumatic stress symptoms, coping self-efficacy, lack of emotional support and social acknowledgement between both victim groups, mixed-effects models were conducted with the same control variables as in the logistic regression analyses.

Differences in incidence (change) between both victim groups

Differences in the incidence of anxiety and depression symptoms, general mental health problems and lack of social support were conducted with similar logistic regression analyses (the incidence of other variables could not be calculated). The incidence was calculated by dividing the number of respondents with post-event symptoms, problems and lack of support among those without pre-event symptoms, problems and lack of support, by the total number of the victim group.

Differences after and before the outbreak among non-victims

To examine the extent to which the prevalence of anxiety and depression symptoms, general mental health problems and lack of social support among non-victims changed after the outbreak (T_4) compared with before the outbreak (T_2), repeated measures multivariate logistic regression analyses were performed. In the analyses, we controlled for T_2 - and T_4 -related demographics and reported other potentially traumatic or SLEs. Mixed-effects models were conducted to examine differences in mean scores of symptoms, problems, and support at T_2 and T_4 on the same (time-dependent) variables.

All analyses were performed with IBM SPSS version 26.

Results

Characteristics of victims and non-victims

The characteristics of the two victim groups, including pre-event are represented in Table 1. Table 1 shows that these two groups differed in gender, number of different VAT events, and the prevalence of general mental health problems at a $P \leq 0.10$ level.

The characteristics of the non-victims at T_2 and T_4 are presented in Table 2. During the 12 months before T_4 , non-victims were significantly less often exposed to SLEs than in the 12 months before T_2 (29.7% v. 24.8%, adjusted odds ratio (aOR) = 0.74,

Table 1 Characteristic victims before and after the coronavirus disease 2019 (COVID-19) outbreak^a

	Victimised in 12 months before		χ^2	d.f.	P
	March 2019 (T ₂) (n = 421)	March 2021 (T ₄) (n = 319)			
Gender					
Men	206 (48.9)	178 (55.8)	3.429	1	0.064
Women	215 (51.1)	141 (44.2)			
Age, years					
18–34	116 (27.6)	102 (32.0)	2.075	3	0.557
35–49	121 (28.7)	87 (27.3)			
50–64	107 (25.4)	80 (25.1)			
65 or older	77 (18.3)	50 (15.7)			
Education level ^b					
Low	94 (22.3)	65 (20.4)	0.628	2	0.731
Medium	147 (34.8)	119 (37.3)			
High	181 (42.9)	135 (42.3)			
Employed					
No	168 (39.9)	134 (42.0)	0.332	1	0.565
Yes	253 (60.1)	185 (58.0)			
Married					
Yes	172 (40.9)	117 (36.7)	1.331	1	0.249
No	249 (59.1)	202 (63.3)			
Physical problems					
Yes	227 (53.8)	181 (56.7)	0.638	1	0.424
No	195 (46.2)	138 (43.3)			
Other PTE/SLEs in past 12 month					
No	202 (47.9)	156 (48.9)	0.078	1	0.780
Yes	220 (52.1)	163 (51.1)			
Period VAT event ^c					
Up to 2 months ago	148 (36.3)	115 (38.3)	0.314	1	0.575
3–12 months	260 (63.7)	185 (61.7)			
Stress during VAT event ^c					
Low	241 (59.2)	161 (53.8)	2.025	1	0.155
High	166 (40.8)	138 (46.2)			
Different types of VATs ^d in past 12 month					
1	350 (83.1)	249 (78.1)	3.035	1	0.081
2 or more	71 (16.9)	70 (21.9)			
VAT in year before					
No	304 (72.0)	213 (66.8)	2.389	1	0.122
Yes	118 (28.0)	106 (33.2)			
Anxiety and depressions symptom year before					
No	308 (73.2)	223 (69.9)	0.948	1	0.330
Yes	113 (26.8)	96 (30.1)			
General mental health problems year before					
No	339 (80.5)	241 (75.5)	2.649	1	0.104
Yes	82 (19.5)	78 (24.5)			
Lack support year before					
No	346 (82.2)	267 (83.7)	0.293	1	0.589
Yes	75 (17.8)	52 (16.3)			

PTE, potentially traumatic event; SLE, stressful life event; VAT, exposure to violence, accidents and/or serious threat.
a. As a result of weighting, the numbers (n) may differ slightly.
b. Low, primary school, intermediate secondary education (USA: junior high school); medium, higher secondary education/preparatory university education (USA: senior high school); intermediate, vocational education (USA: junior college); high, higher vocational education, (USA: college, university), according to education level categories of Statistics Netherlands (CBS).
c. Respondents who were confronted with more than one type of VAT event, were asked to have the most stressful VAT event in the past 12 months in mind when answering the question. Of all 740 VAT victims, 33 answered this question with a theft-related event in mind. They were omitted from these analyses (n^{victims group 1} = 407, n^{victims group 2} = 300).
d. Three types of VATs were distinguished: violence, accidents and serious threats.

95% CI 0.67–0.82, P < 0.001). In addition, the prevalence of physical problems decreased significantly (30.7% v. 29.6%, aOR = 0.992, 95% CI 0.862–0.988, P = 0.020) among non-victims.

Differences between the two victim groups

Table 3 shows that respondents victimised after the outbreak of COVID-19 have a significant higher prevalence (37.6% v. 26.4%)

Table 2 Characteristics non-victims^a

	Non-victims (n = 3245)	
	March 2019 (T ₂), n (%)	March 2021 (T ₄), n (%)
Gender		
Men	1585 (48.8)	1585 (48.8)
Women	1661 (51.2)	1661 (51.2)
Age, years		
18–34	821 (25.3)	706 (21.7)
35–49	758 (23.4)	749 (23.1)
50–64	844 (26.0)	836 (25.8)
65 or older	822 (25.3)	955 (29.4)
Education ^b		
Low	818 (25.2)	781 (24.1)
Medium	1180 (36.4)	1143 (35.2)
High	1248 (38.4)	1321 (40.7)
Employed		
No	1495 (46.1)	1499 (46.2)
Yes	1750 (53.9)	1746 (53.8)
Married		
Yes	1632 (50.3)	1670 (51.5)
No	1613 (49.7)	1575 (48.5)
Physical problems		
Yes	997 (30.7)	960 (29.6)
No	2248 (69.3)	2285 (70.4)
Other PTE/SLE		
No	2281 (70.3)	2440 (75.2)
Yes	964 (29.7)	805 (24.8)

PTE, potentially traumatic event; SLE, stressful life event.
a. As a result of weighting, the numbers (n) may differ slightly.
b. Low, primary school, intermediate secondary education (USA: junior high school); medium, higher secondary education/preparatory university education (USA: senior high school); intermediate, vocational education (USA: junior college); high, higher vocational education, (USA: college, university), according to education level categories of Statistics Netherlands (CBS).

and incidence (13.8% v. 8.8%) of anxiety and depression symptoms, and lower mean scores (i.e. higher levels) of these symptoms (64.1 v. 69.6, Cohens d = 0.30) than respondents victimised before the outbreak, while controlling for gender, the number of different VAT events in the past 12 months and general mental health problems in the year before.

In addition, they also have a significant higher prevalence of general mental health problems (30.4% v. 18.8%) and a significant higher prevalence of probable PTSD (28.1% v. 19.9%). Adults victimised in the year after the outbreak of COVID-19 had also significant higher post-traumatic symptom scores (Cohen’s d = 0.25) than victims before the outbreak. In addition, this group had a significant lower prevalence of high coping self-efficacy levels (81.0% v. 89.4%) than adults victimised before this pandemic, but mean scores did not differ significantly between both groups. No significant differences were found between the two victim groups in lack of emotional support and social acknowledgement.

Differences after and before the outbreak among non-victims

Repeated measures multivariate logistic regression analyses (not shown in table) among the non-victims (n = 3245) revealed a significantly higher prevalence of anxiety and depression symptoms at T₄ (15.0%) than at T₂ (14.5%; aOR = 1.14, 95% CI 1.02–1.26, P = 0.015). No significant (P < 0.05) differences in general mental health problems (9.5% v. 9.2%), or lack of emotional support (8.2% v. 8.7%) were found. Mixed-effects models showed no significant differences in anxiety and depression symptom levels between T₄ and T₂ (T₂ mean 75.6, s.d. = 15.95; T₄ mean 75.40, s.d. = 15.70) and no differences in lack of emotional support levels (T₂ mean 22.1, s.d. = 3.24; T₄ mean 21.9, s.d. = 3.28).

Table 3 Differences between victims before and victims after the coronavirus disease 2019 (COVID-19) outbreak^a

	Victimised in 12 months before		aOR (95% CI)	F (1, 739)	P
	March 2019 (T ₂) (n = 421)	March 2021 (T ₃) (n = 319)			
Anxiety and depression symptoms ^b					
Score, mean (s.d.)	69.6 (19.4)	64.1 (19.8)		11.228	0.001
Prevalence, n (%)	111 (26.4)	120 (37.6)	1.63 (1.15–2.29)		0.005
Incidence, n (%)	37 (8.8)	44 (13.8)	1.64 (1.02–2.61)		0.039
General mental health problems					
Score, mean (s.d.)	n.a.	n.a.			
Prevalence, n (%)	79 (18.8)	97 (30.4)	1.92 (1.26–2.92)		0.002
Incidence, n (%)	31 (7.4)	36 (11.3)	1.56 (0.93–2.61)		0.089
Post-traumatic stress symptoms ^c					
Score, mean (s.d.)	5.6 (6.8)	7.4 (7.9)		7.636	0.006
Prevalence, n (%)	81 (19.9)	84 (28.1)	1.48 (1.02–2.14)		0.040
Incidence, n (%)	n.a.	n.a.			
Coping self-efficacy ^c					
Mean score	39.4 (9.6)	38.1 (9.9)		0.795	0.373
Prevalence, n (%)	364 (89.4)	243 (81.0)	0.55 (0.35–0.87)		0.011
Incidence, n (%)	n.a.	n.a.			
Lack of emotional support					
Score, mean (s.d.)	20.4 (4.0)	20.5 (4.2)		0.633	0.427
Prevalence, n (%)	73 (17.3)	57 (17.9)	0.98 (0.66–1.45)		0.913
Incidence, n (%)	38 (9.0)	28 (8.8)	0.94 (0.56–1.58)		0.822
Lack social recognition ^{c,d}					
Score, mean (s.d.)	10.4 (4.7)	11.2 (4.7)		2.422	0.120
Prevalence, n (%)	37 (9.6)	41 (14.3)	1.44 (0.88–2.35)		0.143
Incidence, n (%)	n.a.	n.a.			

aOR, adjusted odds ratio; n.a., not applicable.

^b As a result of weighting, the numbers (n) may slightly differ. aOR and F-value adjusted for gender, number of different exposure to violence, accidents and/or serious threat (VAT) events in the past 12 months, and general mental health problems in the year before. Incidence is incidence for group 1^{victims} prevalence in March 2019 among those without symptoms, problems and lack support in March 2018, and for group 2^{victims} prevalence in March 2021 among those without symptoms, problems and lack support in March 2020 (according to criteria prevalence).

^c Lower scores indicate higher anxiety and depression symptom/higher lack of emotional support levels.

^d Respondents who were confronted with more than one type of VAT event, were asked to have the most stressful VAT event in the past 12 months in mind when filling in the questionnaire. Of all 740 VAT victims, 33 filled in the PTSD Checklist for DSM-5 /seven-item Coping Self-Efficacy/Social Acknowledgment Questionnaire with a theft-related event in mind. They were omitted from the analyses (n^{victims group 1} = 407, n^{victims group 2} = 300). Including this small group showed almost similar results (see Supplementary Table 1, available at <https://doi.org/10.1192/bjp.2021.226>).

^e n^{victims group 1} = 387, n^{victims group 2} = 287.

Differences between victims and non-victims

The prevalence of moderate–severe anxiety and depression symptom levels, general mental health problems and high lack of emotional support were significantly more prevalent among victims than among non-victims in the same survey, when controlling for demographics and other SLEs (2019: aOR^{anxiety and depression symptoms} = 1.84, 95% CI 1.43–2.37, $P < 0.001$; aOR^{general mental health problems} = 2.01, 95% CI 1.51–2.66, $P < 0.001$; and aOR^{lack of emotional support} = 2.17, 95% CI 1.63–2.90, $P < 0.001$, respectively; 2021: aOR^{anxiety and depression symptoms} = 2.77, 95% CI 2.13–3.60 $P < 0.001$; aOR^{general mental health problems} = 3.34, 95% CI 2.51–4.45, $P < 0.001$; and aOR^{lack of emotional support} = 2.23, 95% CI 1.72–2.90, $P < 0.001$, respectively).

Post hoc control analyses

Respondents who were victimised in both study periods were randomly allocated to group 1^{victims} or to group 2^{victims}. Although both victim groups did not significantly differ in exposure to VATs in the year before, to rule out the possibility that the described results can be attributed to (non-significant) differences in exposure to VATs in the year before (28.0% v. 33.2%) we reanalysed the data by adding this variable to the list of control variables. No relevant differences to the results presented above emerged.

In addition, analyses of the prevalence of specific VAT-related events showed that group 1^{victims} significantly more often ($P < 0.05$) was exposed to traffic accidents (36.1% v. 29.1%) than group 2^{victims}, but that group 2^{victims} was significant more often exposed to physical violence (not by own partner): 17.6% v. 9.7% (see Supplementary Table 2 available at <https://doi.org/10.1192/bjp.2021.226>). There was a statistical trend ($P < 0.10$) that group

2^{victims} were also more often exposed to robberies (6.3% v. 3.3%). However, the fact that group 2^{victims} was more often exposed to these events did not explain the differences in anxiety and depression symptoms, general mental health, PTSD symptomatology and coping self-efficacy. Adding these two variables to the list of control variables did not change results. Both groups did not significantly differ in the distribution of (the most stressful) VAT event when filling in the PCL-5, CSE-7 and SAQ ($\chi^2 = 12.21$, d.f. = 12, $P = 0.429$, see Supplementary Table 3).

Discussion

The general aim of the present study was to examine the extent to which the COVID-19 pandemic and all related disruptions hindered the recovery of victims of potentially traumatic events. For this purpose, we compared the mental health, including PTSD symptomatology, trauma-related coping self-efficacy, and lack of emotional support and social acknowledgement among adults victimised during this pandemic (in the period March 2020–March 2021) with adults victimised before this pandemic (March 2018–March 2019). We focused on VATs.

Findings showed the following clear patterns. First, this pandemic partly hindered the recovery of victims as indicated by a significant higher prevalence and incidence of anxiety and depression symptoms, higher prevalence of general mental health problems, a lower prevalence of high coping self-efficacy and higher post-traumatic symptom scores among adult victims 12 months after the outbreak of COVID-19 compared with adults victimised in the same period before the outbreak. Second, no indications were found that this pandemic had a negative effect on emotional

support and event-related social acknowledgement: both victim groups did not differ in related prevalence, incidence and/or mean scores, although they more often lacked emotional support than non-victims at both March 2019 and March 2021. These findings suggest that the decreased recovery among victims after the outbreak cannot be attributed to a decrease of emotional support, but to other stressors related to this pandemic (such as lockdowns, infection, health risks, unemployment risk).

Although there was no significant difference in the mean scores for anxiety and depression symptoms, the proportion of non-victims scoring above the cut-off score for elevated symptom levels slightly increased between before and after the COVID-19 outbreak (14.5% to 15.0%), whereas the prevalence of physical problems slightly decreased. The finding that such trivial differences gain statistical significance is most probably related to the large sample size ($n = 3245$). Among non-victims, no significant differences between both time periods were found with respect to general mental health problems and lack of emotional support. These results indicate that this pandemic particularly affected victims of VATs. Previous population-based research in the Netherlands found no indications for an increase in the prevalence and incidence of mental health problems among the general population until November–December 2020 compared with pre-pandemic levels, in contrast to an increase in loneliness.^{22,23}

Practical relevance

A large number of COVID-19-related studies are aimed at identifying groups that are at risk for mental health problems because of this pandemic. These insights may guide the development and implementation of programmes and interventions to target these groups at risk so as to prevent or reduce (ongoing) COVID-19-related mental health problems. Victims of VATs are already at risk for mental health problems in 'normal' times, but this study clearly shows that being a victim of these events during this pandemic amplifies this risk. For mental healthcare professionals, general practitioners and victim services, screening and monitoring of victims' mental health are important measures to determine if and which further interventions are needed. Our results indicate that during this pandemic these measures become even more important, also because lower coping self-efficacy levels were observed that increase the risk for mental health problems.^{4,24}

These findings stress the importance of securing the accessibility of mental healthcare services that was hindered during this pandemic because of social distancing measures, for instance, by providing virtual 'walk-in' clinics/victim services.²⁵ Importantly, as no significant differences between the two victim groups in lack of emotional support and social acknowledgement were found, it is not plausible to attribute the amplified risk to diminished social support levels. However, like in pre-pandemic times, victims more often lack emotional support than non-victims. It therefore remains equally important to address the lack of social support among victims.

Strength and limitations

To the best of our knowledge, this is the first comparative longitudinal study assessing the impact of the COVID-19 pandemic on the mental health, coping self-efficacy and social support of victims of VATs. Major strengths are the use of a large longitudinal population-based random sample, the high response rates of each survey, the weighting of data to optimise the representativeness of the results, the use of validated questionnaires, analyses in which differences between both victim groups were controlled for, and the inclusion of non-victims assessed in the same period. However, some limitations need to be clarified.

Our study did not include children and adolescents, we did not examine the effects of this pandemic on respondents who were victimised by earlier potentially traumatic events such as adverse childhood experiences, and no clinical interviews were conducted to examine mental disorders among victims and non-victims. We have no information about the frequency of events. Given the aim of the present study, we did not focus on acute, life-threatening and catastrophic COVID-19-related events.^{26,27} In addition, we did not examine other relevant aspects of mental health such as burnout.

The main conclusions that can be drawn from the results is that adults victimised by VATs in the year following the outbreak of COVID-19 more often have post-traumatic stress symptoms, anxiety and depression symptoms, and general mental health problems, and had lower coping self-efficacy levels than those victimised before the outbreak.

Peter G. van der Velden , Centerdata, the Netherlands and Tilburg University's Network on Health and Labor (NETHLAB), Tilburg University, the Netherlands; **Carlo Contino**, Fonds Slachtofferhulp, the Netherlands; **Marcel Das**, Centerdata, the Netherlands and Tilburg School of Economics and Management, Tilburg University, the Netherlands; **Joost Leenen**, Centerdata, the Netherlands; **Lutz Wittmann**, International Psychoanalytic University Berlin, Germany

Correspondence: Peter G. van der Velden. Email: pg.vandervelden@tilburguniversity.edu

First received 3 Oct 2021, final revision 8 Dec 2021, accepted 21 Dec 2021

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1192/bjp.2021.226>

Data availability

The data of the surveys in 2018, 2019, 2020 and 2021 of the VICTIMS study are, after registration, freely available from the LISS dataarchive (see <https://www.dataarchive.lissdata.nl>, in English, assembled study 208) for researchers, like all other studies conducted with the LISS panel. The data of the survey conducted in 2021 will be added to the data archive soon.

Author contributions

P.G.v.d.V. and C.C. conceived and designed the study in collaboration with M.D. and L.W. The data analyses was performed by P.G.v.d.V. in cooperation with M.D. and L.W. All authors contributed to the data interpretation. P.G.v.d.V. wrote the first draft. All authors critically revised the first draft, and all authors approved the final version of the article to be published.

Funding

This study is funded by Fonds Slachtofferhulp, the Netherlands (50006/VICTIMS).

Declaration of interest

None.

References

- van der Velden PG, Komproe I, Contino C, de Bruijne M, Kleber RJ, Das M, et al. Which groups affected by potentially traumatic events (PTEs) are most at risk for a lack of social support? A prospective population-based study on the 12-month prevalence of PTEs and risk factors for a lack of post-event social support. *PLoS ONE* 2020; **15**: e0232477.
- Koenen KC, Ratanatharathorn A, Ng L, McLaughlin KA, Bromet EJ, Stein DJ, et al. Posttraumatic stress disorder in the World Mental Health Surveys. *Psychol Med* 2017; **47**: 2260–74.
- Benight CC, Bandura A. Social cognitive theory of posttraumatic recovery: the role of perceived self-efficacy. *Behav Res Ther* 2004; **42**: 1129–48.

- 4 Luszczyńska A, Benight CC, Cieslak R. Self-efficacy and health-related outcomes of collective trauma: a systematic review. *Euro Psychologist* 2009; **14**: 51–62.
- 5 Kaniasty K, Norris FH. Longitudinal linkages between perceived social support and posttraumatic stress symptoms: sequential roles of social causation and social selection. *J Trauma Stress* 2008; **21**: 274–81.
- 6 van der Velden PG, Contino C, Marchand M, Das M, Schut H. Does pre-event lack of emotional support increase the risk of post-event PTSD, anxiety, depression symptoms and lack of support? A comparative population-based study among victims of threat and violence. *J Anxiety Disord* 2020; **75**: 102269.
- 7 Robinson E, Sutin AR, Daly M, Jones A. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic. *J Affect Disord* 2022; **296**: 567–76.
- 8 Hobfoll SE. Conservation of resources. A new attempt at conceptualizing stress. *Am Psychol* 1989; **44**: 513–24.
- 9 Hobfoll SE. Social and psychological resources and adaptation. *Rev Gen Psychol* 2002; **6**: 307–24.
- 10 Scherpenzeel A, Das M. True longitudinal and probability based internet panels: evidence from The Netherlands. In *Social and Behavioral Research and the Internet: Advances in Applied Methods and Research Strategies* (eds M Das, P Ester and L Kaczmirek): 77–104. Taylor & Francis, 2011.
- 11 Ware JE, Sherbourne CD. The MOS 36-item short-form health survey, SF-36: I. Conceptual framework and item selection. *Med Care* 1992; **30**: 473–83.
- 12 Means-Christensen AJ, Arnau RC, Tonidandel AM, Bramson R, Meagher MW. An efficient method of identifying major depression and panic disorder in primary care. *J Behav Med* 2005; **28**: 565–572.
- 13 Driessen M. *Een beschrijving van de MHI-5 in De Gezondheidsmodule van het Permanent Onderzoek Leefsituatie [A Description of the MHI-5 in the Health Module of Permanent Research of Living Conditions, POLS]*. Statistics Netherlands, 2011.
- 14 Price M, Szafranski DD, van Stolk-Cooke K, Gros DF. Investigation of abbreviated 4 and 8 item versions of the PTSD Checklist 5. *Psychiatry Res* 2016; **239**: 124–30.
- 15 Weathers FW. Posttraumatic stress disorder checklist. In *Encyclopedia of Psychological Trauma* (eds G Reyes and JD Elhai): 491–94. Wiley, 2008.
- 16 Pereira-Lima K, Loureiro SR, Bolsoni LM, Apolinario da Silva TD, Osório FL. Psychometric properties and diagnostic utility of a Brazilian version of the PCL-5 (complete and abbreviated versions). *Eur J Psychotraumatol* 2019; **10**: 1581020.
- 17 van der Velden PG, Das M, Contino C, van der Knaap LM. From health to financial problems: multiproblems among victims of partner and non-partner physical violence, and matched nonvictims. *J Interpers Violence* 2019; **5**: 886260519885915.
- 18 Bosmans MWG, Komproe IH, Van der Knaap LM, Benight CC, van der Velden PG. Assessing perceived ability to cope with trauma: a multigroup validity study of a trauma-related coping self-efficacy scale. *Eur J Psychol Assess* 2017; **33**: 55–64.
- 19 Maercker A, Müller J. Social acknowledgment as a victim or survivor: a scale to measure a recovery factor of PTSD. *J Trauma Stress* 2004; **17**: 345–51.
- 20 Bridges KR, Sanderman R, van Sonderen E. An English language version of the social support list: preliminary reliability. *Psychol Rep* 2002; **90**: 1055–58.
- 21 van Sonderen E. *Het meten van Sociale Steun met de Sociale Steun Lijst - Interacties SSL-I en Sociale Steun Lijst - Discrepanties, SSL-D: een handleiding (Tweede herziene druk) [The measurement of social support with the Social Support List - Interactions (SSL-I) and Discrepancies (SSL-D)]*. Research Institute SHARE RUG, 2012.
- 22 van der Velden PG, Hyland P, Contino C, von Gaudecker HM, Muffels R, Das M. Anxiety and depression symptoms, the recovery from symptoms, and loneliness before and after the COVID-19 outbreak among the general population. Findings from a Dutch population-based longitudinal study. *PLoS ONE* 2021; **16**: e0245057.
- 23 van der Velden PG, Marchand M, Das M, Muffels R, Bosmans M. The prevalence, incidence and risk factors of mental health problems and mental health services use before and 9 months after the COVID-19 outbreak among the general Dutch population. A 3-wave prospective study. *medRxiv* [Preprint] 2021. Available from: <https://doi.org/10.1101/2021.02.27.21251952>.
- 24 Bosmans MWG, van der Velden PG. Cross-lagged associations between post-traumatic stress symptoms and coping self-efficacy in long-term recovery: a four-wave comparative study. *Soc Sci Med* 2017; **193**: 33–40.
- 25 Matthewman S, Spencer S, Lavergne MR, McCracken RK, Hedden L. An environmental scan of virtual “walk-in” clinics in Canada: comparative study. *J Med Internet Res* 2021; **23**: e27259.
- 26 Norrholm SD, Zalta A, Zoellner L, Powers A, Tull MT, Reist C, et al. Does COVID-19 count? Defining Criterion A trauma for diagnosing PTSD during a global crisis. *Depress Anxiety* 2021; **38**: 882–85.
- 27 Bridgland VME, Moeck EK, Green DM, Swain TL, Nayda DM, Matson LA, et al. Why the COVID-19 pandemic is a traumatic stressor. *PLoS ONE* 2021; **16**: e0240146.

